2025/08/29 07:36 1/1 Ideas

Ideas

- Airway
 - Hyperangulated VL vs Standard DL in Surgical Patients
 - Finding: First-pass intubation failure: hyperangulated VL 1.7% vs DL 7.6%
 - **Significance**: More evidence VL is better as first-line?
 - Mac Geometry VL vs Mac DL
 - Finding: VL had better first-pass success than DL
- CARDS
 - IV vs IO for OHCA
 - Finding: IV assoc with better survival and neuro fav outcomes
 - MINT: Restrictive or Liberal Transfusion Strategy in Myocardial Infarction and Anemia
 - **Finding**: In patients with acute myocardial infarction and anemia, a liberal transfusion strategy did not significantly reduce the risk of recurrent myocardial infarction or death at 30 days.
- CCM
 - Pantoprazole for MV GI bleed ppx
 - Finding: reduces GIB, no mortality benefit
 - Calcium for hyperK
 - Finding: not via membrane stabilization
- MCS
 - DANGER-SHOCK
 - Finding: Impella CP reduces mortality
 - Significance: First trial to show mortality benefit from MCS
- PULM
 - brensocatib for bronchiectasis
 - Depemokimab
 - Inebilizumab for Treatment of IgG4-Related Disease
- GI
- Tranexamic acid in upper gastrointestinal bleed in patients with cirrhosis
 - Finding: in Child-Pugh B or C with UGIB, TXA improves bleeding control vs placebo
- Trauma
 - UK-REBOA
 - **Finding**: In trauma patients with exsanguinating hemorrhage, a strategy of REBOA and standard care in the emergency department does not reduce, and may increase, mortality compared with standard care alone.

From:

https://ewrobbins.com/ - ewrobbins.com

Permanent link:

https://ewrobbins.com/doku.php?id=jc

Last update: 2025/04/24 13:52

