Ideas

- DANGER-SHOCK
 - Finding: Impella CP reduces mortality
 - $\,\circ\,$ Significance: First trial to show mortality benefit from MCS
- Hyperangulated VL vs Standard DL in Surgical Patients
 - Finding: First-pass intubation failure: hyperangulated VL 1.7% vs DL 7.6%
 - Significance: More evidence VL is better as first-line?
- Tranexamic acid in upper gastrointestinal bleed in patients with cirrhosis

• Finding: in Child-Pugh B or C with UGIB, TXA improves bleeding control vs placebo

- UK-REBOA
 - **Finding**: In trauma patients with exsanguinating hemorrhage, a strategy of REBOA and standard care in the emergency department does not reduce, and may increase, mortality compared with standard care alone.
- MINT: Restrictive or Liberal Transfusion Strategy in Myocardial Infarction and Anemia
 - Finding: In patients with acute myocardial infarction and anemia, a liberal transfusion strategy did not significantly reduce the risk of recurrent myocardial infarction or death at 30 days.
- Mac Geometry VL vs Mac DL
 - $\circ\,$ Finding: VL had better first-pass success than DL

From: https://ewrobbins.com/ - ewrobbins.com

Permanent link: https://ewrobbins.com/doku.php?id=jc&rev=1715709377



Last update: 2024/05/14 17:56

1/1