# SSc-ILD

- GAP score
- ILD-RISC score
  - Outcome to predict: the presence of ILD on HRCT
  - score: a probability (0 to 1.0)
  - ILD-RISC score >= 0.3 had sensitivity 85.6% and specificity 53.6%
  - items:
    - ☐ age
    - ☐ ever had digital ulcers (OR 2.06)
    - ☐ DLCO
    - ☐ FVC
    - ☐ antibody (topo I: OR 2.38, PM/SCL: 2.56, RNAPIII: 1.41)

# IIM-ILD

- FLAIR Model

| item | categorical_value | score |
|---|---|---|
| ferritin | >= 636 ng/mL | TRUE: 2, FALSE: 0 |
| LDH | >= 355 U/L | TRUE: 2, FALSE: 0 |
| anti-MDA5 antibody | >= 636 ng/mL | negative, 0; +, 2; ++, 3; +++, 4 |
| high-resolution CT imaging score (as described by [https://doi.org/10.1164/rccm.2106157]) | >= 133 | TRUE: 3, FALSE: 0 |

From:
https://ewrobbins.com/ - **ewrobbins.com**

Permanent link:
**https://ewrobbins.com/doku.php?id=resources:pulm:ctd-ild&rev=1793129546**

Last update: **2026/10/27 19:32**