SSc-ILD

- GAP score
- ILD-RISC score
  - Outcome to predict: the presence of ILD on HRCT
  - score: a probability (0 to 1.0)
  - ILD-RISC score >= 0.3 had sensitivity 85.6% and specificity 53.6%
  - items:
    - ☐ age
    - ☐ ever had digital ulcers (OR 2.06)
    - ☐ DLCO
    - ☐ FVC
    - ☐ antibody (topo I: OR 2.38, PM/SCL: 2.56, RNAPIII: 1.41)

IIM-ILD

- FLAIR Model
  - bottom line: MDA5 is bad, high inflammatory markers are bad, l

item	categorical_value	score
ferritin	>= 636 ng/mL	TRUE: 2, FALSE: 0
LDH	>= 355 U/L	TRUE: 2, FALSE: 0
anti-MDA5 antibody	>= 636 ng/mL	negative, 0; +, 2; ++, 3; +++, 4
high-resolution CT imaging score (as described by https://doi.org/10.1164/rccm.2106157)	>= 133	TRUE: 3, FALSE: 0

- Scoring
  - low, 0 to 4
  - medium, 5 to 9
  - high, 10 to 13

From:
https://ewrobbins.com/ - **ewrobbins.com**

Permanent link:
https://ewrobbins.com/doku.php?id=resources:pulm:ctd-ild&rev=1793130497

Last update: **2026/10/27 19:48**